

Your article (Vol. 24, No. 15, mcb0311-04) from Molecular and Cellular Biology is available for download
=====

Molecular and Cellular Biology Published by the American Society for Microbiology

Dear Author,

Please refer to the following URL:

<http://rapidproof.cadmus.com/RapidProof/retrieval/index.jsp>

Login: your e-mail address

Password: ----

The site contains 1 file. You will need to have Adobe Acrobat? Reader software to read these files. This is free software and is available for user downloading at
<http://www.adobe.com/products/acrobat/readstep.html>.

This file contains:

Instructions for proof handling

Proofreading marks guide

Reprint/page charge form

A copy of the page proofs for your article

A query page (if applicable)

After printing the PDF file, please read the page proofs carefully and:

- 1) indicate changes or corrections on the printed page proofs; DO NOT EDIT OR ALTER THE PDF FILE IN ANY WAY;
- 2) answer all queries (footnotes A,B,C, etc.) on the last page of the PDF proof;
- 3) proofread any tables and equations carefully;
- 4) check that any Greek, especially μ , has translated correctly;
- 5) sign and date signature block on the first page of the proofs.

Within 48 hours, please mail the following to the ASM Journals Department at the address given below:

- 1) Original PDF set of page proofs, with your corrections clearly marked and signature provided on the first page. Corrections from all authors must be incorporated in one set of page proofs. Corrected proofs should be mailed or sent via a courier such as FedEx. Faxing is not recommended; ASM will not be responsible for errors caused by poor-quality faxes. DO NOT SEND THE PROOFS AS AN E-MAIL ATTACHMENT.
- 2) Reprint/page charge form.

If you have any problems with your proofs or questions regarding changes you would like to make, please contact me. PLEASE ALWAYS INCLUDE YOUR ARTICLE NO. (mcb0311-04) WITH ALL CORRESPONDENCE.

If you have problems accessing or viewing your PDF proofs, please contact Katie Gay of Cadmus Professional Communications at 804-261-3155 or Email: gayk@cadmus.com.

The proof contains 4 pages.

Sincerely,

Becky Zwadyk
Production Editor
Molecular and Cellular Biology

Journals Department
American Society for Microbiology
1752 N St., N.W.
Washington, DC 20036-2904
Tel: 202-942-9214
Fax: 202-942-9355
Email: bzwadyk@asmusa.org

INSTRUCTIONS

Mark all corrections *on* the page proofs that you print from the PDF file.

Answer all queries.

Check the wording of the running heads. Note that the page numbers on the proofs are for easy reference only; they are not the actual page numbers that will be used for the printed article.

If there are illustrations, see that each is numbered correctly, is matched with the appropriate legend, and is correctly oriented.

Sign your name on the first page of the proofs in the block provided.

Send the corrected proofs and all 5 pages of the reprint order form to the ASM Journals Department *by mail, express mail, or courier* within 48 h after you receive them. (Corrections received by ASM after the date indicated in the acceptance letter might not be incorporated in the published article.)

ILLUSTRATIONS

Both black-and-white and color images represent samples of the graphic file that will be used for printing. Final output will be a higher-resolution version than what appears in the PDF.

ALTERATIONS

You are provided with a proof of your paper primarily to check the accuracy of the type. *The final responsibility for correcting all errors is yours.*

However, the proof stage is *not* the time for revision, rewriting, rephrasing, addition of more recent material, or any other significant change from the final edited manuscript. That is, the manuscript that was approved by the editor should be the one printed; there should be no major additions or deletions. In case of *essential* new information, you may send a *short* “Addendum in Proof,” provided that the editor has given his or her consent. If references to unpublished data or personal communications are added, it is expected that written assurance granting permission for the citation will be included.

This journal is published on a rapid schedule. Major additions or deletions of material cannot be considered, except under extraordinary conditions, and the cost of such alterations may be billed to the author.

NOTE: It is recommended that you make a copy of the corrected proofs for your file.

Proofreaders' Marks and How to Use Them

Use of proofreaders' marks requires a double marking—a mark in the body of the text at the point of correction and the corresponding correction in the margin. A correction marked only in the text and not in the margin will be missed by the printer.

The following is a list of the proofreaders' marks in general use.

PUNCTUATION		SPACING	
⊙	Period	#	Insert space
↗	Comma	eq #	Equalize space
⊙	Colon	⌋	Close up
↗	Semicolon	STYLE OF TYPE	
↘	Apostrophe	wf	Wrong font
↘	Open quotes	lc	Lower case
↘	Close quotes	cap	Capitalize
=	Hyphen	c lc	Initial cap, then lower case
—	Dash (show length)	sc	Small capitals
()	Parentheses	c sc	Initial cap, then small caps
DELETE AND INSERT		rom	Set in roman
→	Delete	ital	Set in italics
↻	Delete and close up	lf	Set in light face
see act copy	Insert omitted matter	bf	Set in bold face
stet	Let it stand	3	Superior character
PARAGRAPHING		3	Inferior character
¶	Paragraph	MISCELLANEOUS	
fl ¶	Flush paragraph	X	Broken type
¶ ¶	Indent (show no. of ems)	9	Invert
run in	Run in	↓	Push down
POSITION		sp	Spell out
☐ ☐	Move right or left	/	Shilling mark (slash)
☐ ☐	Raise, lower	⋯	Ellipsis
ctr	Center	see l/o	See layout
fl L fl R	Flush left, right	? query	Query
=	Align horizontally		
	Align vertically		
tr	Transpose		
tr #	Transpose space		

MOLECULAR AND CELLULAR BIOLOGY

PROOFREADING

Read, correct, sign, and return this set of proofs *within 48 hours*. Mail corrected proofs and this form to:

**Journals Department
American Society for Microbiology
1752 N St., NW
Washington, DC 20036-2904
USA**

IMPORTANT

This form must be completed and returned with the corrected page proofs.

To guarantee more efficient service, we request that all information be typed or clearly printed on this form. Illegible handwriting may result in shipment to the wrong address. If any such orders have to be reshipped, the additional expense will be charged to the customer.

Questions?

About this reprint/page charge form

Contact: Mary Hill, Finance Dept.
Phone, (202) 942-9256
Fax, (202) 942-9370
E-mail, mhill@asmusa.org

About your page proofs

Contact: MCB production editor
Phone, (202) 737-3600
FAX, (202) 942-9355

MOLECULAR AND CELLULAR BIOLOGY

Manuscript control no. _____ Month of publication _____
 Volume no.: 24 Issue no. _____ Year of publication 2004
 Authors _____
 Title _____

IMPORTANT: *Corresponding* authors who are ASM members may take advantage of the reduced reprint/offprint charges and page charges shown in this order form. The corresponding author must be a member in the year in which the article is being published.

MEMBER Corresponding Author

Name: _____

Member no.: _____
 (*or date membership application submitted*)

Black-and-White Reprint Prices*

Pages	Price					
	100 copies	200 copies	300 copies	400 copies	500 copies	Additional 100's
4	\$ 95	\$106	\$119	\$130	\$141	\$13
8	191	213	233	255	275	40
12	282	315	345	378	411	55
16	370	414	458	499	543	66
20	458	513	570	625	680	77

Reprints are supplied only in multiples of 100. To expedite delivery, all reprints are handled in the same way at the same time. Orders for covers or "special printing" cannot be accepted. Orders received after publication of the issue may be subject to a 20% surcharge.

How to determine the cost of black-and-white reprints:

(i) Count the total number of pages; a fraction of a page is considered to be one page. (Note: The proofs sent by the printer are page proofs, not galley proofs.)

(ii) Convert the figure to the next larger whole even number shown in the reprint price table and apply the appropriate rate.

*Price includes cost of UPS shipping for U.S. orders. For non-U.S. orders, see shipping charge chart on next page.

Page Charges

Count the total number of pages, as indicated in step *i* (above), *multiply* each of the first 9 pages by **\$63** and each additional page by **\$86**, and add the products. The sum is the article page charge for member corresponding authors. (Note: *Minireviews, commentaries, and regular letters to the editor are not subject to page charges.*)

NONMEMBER Corresponding Author

Black-and-White Reprint Prices*

Pages	Prices					
	100 copies	200 copies	300 copies	400 copies	500 copies	Additional 100's
4	\$108	\$120	\$135	\$148	\$160	\$15
8	218	243	265	290	313	45
12	320	358	393	430	468	63
16	420	470	520	568	618	75
20	520	583	648	710	773	88

Reprints are supplied only in multiples of 100. To expedite delivery, all reprints are handled in the same way at the same time. Orders for covers or "special printing" cannot be accepted. Orders received after publication of the issue may be subject to a 20% surcharge.

How to determine the cost of black-and-white reprints:

(i) Count the total number of pages; a fraction of a page is considered to be one page. (Note: The proofs sent by the printer are page proofs, not galley proofs.)

(ii) Convert the figure to the next larger whole even number shown in the reprint price table and apply the appropriate rate.

*Price includes cost of UPS shipping for U.S. orders. For non-U.S. orders, see shipping charge chart on next page.

Page Charges

Count the total number of pages, as indicated in step *i* (above), *multiply* each of the first 9 pages by **\$72** and each additional page by **\$99**, and add the products. The sum is the article page charge for nonmember corresponding authors. (Note: *Minireviews, commentaries, and regular letters to the editor are not subject to page charges.*)

BLACK-AND-WHITE Reprint & Page Charges

Shipping Information

Reprints are shipped about 4 weeks after the issue is printed.

U.S. orders are shipped via UPS.

Foreign orders are shipped via an expedited, traceable system.

Charges for Shipping to Foreign Countries

Pages	Number of Copies					
	100 copies	200 copies	300 copies	400 copies	500 copies	Additional 100's
4	\$25	\$ 35	\$ 49	\$ 60	\$ 74	\$15
8	35	60	84	109	134	25
12	49	84	124	158	199	37
16	60	109	158	209	258	50
20	74	134	199	258	318	62

Determine the number of pages as described on the previous page.

Shipping Instructions

(Only one shipping address allowed per order. A street address is required; **we are unable to ship to a P.O. box.**)

Name _____

Address _____

City _____ State _____ Zip _____

Country _____

Fax _____

Charges to the Corresponding Author

No. of Pages: _____

Page Charges (*minireviews, commentaries, and comment letters to the editor are not subject to page charges*):

Member corresponding author page charges _____

Nonmember corresponding author page charges _____

Reprint Charges:

100 free reprints (*minireviews and commentaries only*) _____

Number of reprints being purchased _____

* Member corresponding author reprint charge _____

* Nonmember corresponding author reprint charge _____

Foreign Shipping Charge for Purchased Reprints: _____

TOTAL COST: _____

* Canadian residents must add 7% GST (plus 8% HST where applicable) for reprints.

COLOR Offprint & Page Charges

Shipping Information

Offprints are shipped about 4 weeks after the issue is printed.

U.S. orders are shipped via UPS.

Foreign orders are shipped via an expedited, traceable system.

Charges for Shipping to Foreign Countries

Pages	Charge					
	100 copies	200 copies	300 copies	400 copies	500 copies	Additional 100's
4	\$ 25	\$ 35	\$ 49	\$ 60	\$ 74	\$15
8	35	60	84	109	134	25
12	49	84	124	158	199	37
16	60	109	158	209	258	50
20	74	134	199	258	318	62

Shipping Instructions

(Only one shipping address allowed per order. A street address is required; **we are unable to ship to a P.O. box.**)

Name _____

Address _____

City _____ State _____ Zip _____

Country _____

Fax _____

Charges to the Corresponding Author

No. of Pages: _____

Page Charges (*minireviews, commentaries, and comment letters to the editor are not subject to page charges*):

Member corresponding author page charges _____

Nonmember corresponding author page charges _____

Color Figure Charges (*per cost estimate sent to you by the ASM production editor*):

Member corresponding author color charge: _____

Nonmember corresponding author color charge: _____

Color Offprint Charges (*per cost estimate sent to you by the ASM production editor*):

100 free offprints (*minireviews and commentaries only*) _____

Number of offprints being purchased _____

* Member corresponding author offprint charge _____

* Nonmember corresponding author offprint charge _____

Foreign Shipping Charge for Purchased Color Offprints:

TOTAL COST: _____

* Canadian residents must add 7% GST (plus 8% HST where applicable) for offprints.

Payment by Credit Card*

___ MasterCard ___ VISA ___ AmEx ___ Eurocard

Card number _____

Exp. date _____ Phone no. _____

Name as it appears on card _____

Cardholder signature _____

Cardholder address _____

City _____ State _____ Zip _____

Country _____

* Credit card will be charged in publication month.

Invoicing Instructions

- ☐ Order to be paid by individuals **will not be processed without the signature** of the individual who is ordering the reprints.

Signature (reprints) _____

Signature (page charges) _____

- ☐ Order to be paid by universities, hospitals, corporations, government agencies, or other institutions **will not be processed without a purchase order.**

Purchase order number _____ is

☐ attached

☐ to follow separately§

The purchase order must include the name of the journal, manuscript control number shown above, volume number, issue number, and month and year of publication. **Make the purchase order out to: American Society for Microbiology.**

§ For institutions, a 7-day grace period for return of this form will be granted if necessary for acquisition of a purchase order.

Billing Information

If payment does not accompany this form, give the name and address of the individual or institution to be billed for the reprint/page charges.

Name _____

Address _____

City _____ State _____ Zip _____

Country _____

Telomere-Associated Protein TIN2 Is Essential for Early Embryonic Development through a Telomerase-Independent Pathway

Y. Jeffrey Chiang,^{1*} Sahn-Ho Kim,² Lino Tessarollo,³ Judith Campisi,² and Richard J. Hodes^{1,4}

Experimental Immunology Branch, National Cancer Institute,¹ and National Institute on Aging,⁴ National Institutes of Health, Bethesda, Maryland 20892; Life Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720²; and Center for Cancer Research, Mouse Cancer Genetics Program, National Cancer Institute, Frederick, Maryland 21702³

Received 19 February 2004/Returned for modification 1 May 2004/Accepted 11 May 2004

TIN2 is a negative regulator of telomere elongation that interacts with telomeric DNA repeat binding factor 1 (TRF1) and affects telomere length by a telomerase-dependent mechanism. Here we show that inactivation of the mouse TRF1-interacting protein 2 (TIN2) gene results in early embryonic lethality. We further observed that the embryonic lethality of TIN2 mutant mice was not affected by inactivation of the telomerase reverse transcriptase gene, indicating that embryonic lethality is not the result of telomerase-dependent changes in telomere length or function. Our findings suggest that TIN2 has a role independent of telomere length regulation that is essential for embryonic development and cell viability.

Fn* Telomeres are complex structures at chromosome ends that function to prevent chromosome fusions and genomic instability (6). Mammalian telomeres consist of double-stranded TTAGGG repeats and associated proteins. A number of these telomere-associated proteins have been identified, including telomeric repeat binding factors 1 and 2 (TRF1 and TRF2) (5), TRF1-interacting protein 2 (TIN2) (12), tankyrase 1 (19) and 2 (9), protein interacting with NIMA-interacting factor 1 (22), ATM (13), Ku (8), Rap1 (14), DNA-PK (7), and POT1 (1, 15). TRF1 is a ubiquitously expressed protein that binds to double-stranded telomeric DNA repeats (2, 18). The acidic amino terminus of TRF1 contains a conserved homodimerization motif, and the carboxyl terminus of TRF1 has a DNA-binding domain homologous to Myb oncoprotein (4, 5). TRF1 acts as a negative regulator of telomere length maintenance, with overexpression of TRF1 resulting in telomere shortening and expression of dominant-negative (DN) TRF1 causing telomerase-dependent telomere elongation (2, 21).

TIN2 is a protein that interacts directly with TRF1 and is an essential mediator of TRF1 activity (11, 12). In the presence of telomerase activity, overexpression of TIN2 inhibits telomere elongation in human cell lines, whereas expression of DN N-terminal deletion mutants of TIN2 enhances telomere elongation (12). The TIN2 protein contains N-terminal basic and acidic regions, a central TRF1-binding domain, and a C-terminal region. The basic and acidic regions are required for the regulation of TRF1 activity. The TRF1-binding domain of TIN2 associates with the TRF1 homodimerization domain, providing for the recruitment of TIN2 to the telomere (12). Kim et al. (11) have suggested that binding of wild-type TIN2 induces changes in TRF1 conformation that in turn favor a telomeric structure which is inaccessible to telomerase, thus preventing telomerase-mediated telomere elongation. The ab-

sence of TIN2 would conversely favor telomerase accessibility and telomere elongation.

The physiological role of TIN2 during in vivo development and in normal cell function has not previously been assessed. To better understand the in vivo function of TIN2, we have therefore studied the effect of TIN2 mutation on mouse development by using gene targeting technology. We found that TIN2 deletion results in early embryonic lethality through a mechanism that is independent of telomerase function.

MATERIALS AND METHODS

Targeting vector, electroporation, and selection. For the TIN2 gene, a replacement type targeting vector was constructed from an 8.0-kb 129/Sv (Stratagene) mouse genomic fragment containing the TIN2 coding sequence (Fig. 1A). The *neo* gene with the phosphoglycerate kinase 1 promoter and the bovine growth hormone polyadenylation sequence (pGKneobpA) was employed as a positive selectable marker; the pGK-thymidine kinase cassette was used as a negative selectable marker. Electroporation and selection were performed with the C17 embryonic stem (ES) cell line as described elsewhere (20). DNA from G418- and 1-(2-deoxy-2-fluoro- β -D-arabinofuranosyl)-5-iodouracil (FIAU)-resistant ES clones was screened by diagnostic *NotI* restriction enzyme digestion and with a 3' probe external to the targeting vector sequence, as shown in Fig. 1A. Recombinant clones containing the predicted 13-kb rearranged band were obtained at a frequency of 1/15.

Generation of mutant mice. Two independent ES cell clones targeted for the TIN2 recombination were injected into C57BL/6 blastocysts and generated chimeras that transmitted the mutated allele to progeny (3). Mice were bred in a specific-pathogen-free facility with food and water ad libitum.

Mice. TIN2 heterozygotes were maintained on a *Mus musculus* 129 background or backcrossed to an *M. musculus* C57BL/6 (B6) background. TIN2^{+/-} TERT^{-/-} mice were generated by breeding TIN2^{+/-} mice with TERT^{-/-} mice. The generation and characterization of mTERT knockout mice is described elsewhere (33). Chiang et al., submitted for publication). All animals were housed at Bioqual (Rockville, Md.).

DNA isolation. ES cells and mouse tail preparations were lysed in 500 μ l of DNA lysis buffer containing 100 mM Tris-HCl (pH 8.0), 200 mM NaCl, 10 mM EDTA, 2% sodium dodecyl sulfate, and 50 μ g of proteinase K at 55°C overnight. Lysates were extracted with 500 μ l of phenol. DNA was collected from the upper phase by ethanol precipitation and centrifugation at 12,000 rpm and dissolved in 200 μ l of Tris-EDTA buffer.

Southern blot analysis. For Southern blot analysis, 50 μ g of DNA was digested with *NotI*, electrophoresed on a 0.7% agarose gel, transferred to a nylon membrane, and probed with probe A (Fig. 1B).

PCR genotyping. DNA was prepared as described above, and PCR was performed. Oligonucleotides used for genotyping of TIN2 were NEO1 (5'-TTC

* Corresponding author. Mailing address: Experimental Immunology Branch, National Cancer Institute, National Institutes of Health, Building 10, 4B36, 9000 Rockville Pike, Bethesda, MD 20892. Phone: (301) 496-1376. Fax: (301) 496-0887. E-mail: chiangj@mail.nih.gov.




FIG. 1. Generation of TIN2-deficient mice by gene targeting. (A) Gene targeting strategy and restriction map of the gene. The open boxes represent the exons of the TIN2 gene. A neomycin resistance gene was inserted into exon 1 5' of the translation initiation codon ATG. (B) Southern blot analysis of mouse tail DNA. The 8.5- and 13-kb bands represent the germ line and targeted alleles, respectively. (C) PCR analysis for the TIN2 genotype. The 220- and 270-bp PCR products represent the wild-type and targeted alleles, respectively.

TGG ATT CAT CGA CTG TG-3'), TINF1 (5'-TAC GTA CTC TTG CGT TTC CCT C-3'), and TINR1 (5'-TTC TCT CAT CTG CAC CGG AAG G-3'). The amplification reaction was carried out in a 25- μ l volume containing 1 to 100 ng of DNA, 25 pmol of each oligonucleotide, and 12.5 μ l of Supremix buffer (Invitrogen). PCR conditions were 35 rounds of 95°C for 30 s, 58°C for 30 s, and 72°C for 1 min. PCR products were revealed by agarose gel electrophoresis and ethidium bromide staining. The 220-bp band corresponds to the wild type, and the 270-bp band corresponds to knockout TIN2 (Fig. 1C).

Genotyping of embryos at day 7.5 and 10.5. Timed pregnant mice were prepared by intercrossing TIN2^{+/-} mice. Embryos were isolated, digested, and genotyped by PCR as described above.

In vitro culture and genotyping of blastocysts. Three- to four-day-old embryos from TIN2^{+/-} crosses were isolated and cultured at 37°C in ES cell RPMI medium containing 15% fetal bovine serum and 100 U of LIF/ml (16, 17). Cells were photographed and then digested in 100 μ l of DNA lysis buffer as described above. DNA was precipitated, centrifuged, and dissolved in 10 μ l of H₂O. Genotyping was carried out as described above except that PCR conditions were 5 rounds of 95°C for 30 s, 49°C for 30 s, and 72°C for 1 min; 5 rounds of 95°C for 30 s, 52°C for 30 s, and 72°C for 1 min; 5 rounds of 95°C for 30 s, 55°C for 30 s, and 72°C for 1 min; and 35 rounds of 95°C for 30 s, 58°C for 30 s, and 72°C for 1 min.

RESULTS

Disruption of the TIN2 gene results in early embryonic lethality. In vitro studies have demonstrated that TIN2 inter-

TABLE 1. There are no TIN2^{-/-} offspring from TIN2^{+/-} crosses **AQ: D**

Result	No. of TIN2 ^{+/-} \times TIN2 ^{+/-} offspring		
	TIN2 ^{+/+}	TIN2 ^{+/-}	TIN2 ^{-/-}
Expected	59	118	59
Observed	80	157	0

acts with TRF1 and is an essential mediator of TRF1 activity in telomerase-dependent telomere length regulation. However, little is known about the in vivo function of TIN2 in mammalian cells and during development. To better understand the physiological roles of TIN2, mouse gene targeting technology was used. It has been shown that the B6 mouse TIN2 gene is located on chromosome 14 (GenBank), is approximately 3.98 kb in length, and consists of 9 exons. Before 129/SvJ strain mouse genomic DNA was used to make a gene targeting construct, the TIN2 gene was mapped with 12 common DNA restriction enzymes. The mapping results showed that the 129/SvJ strain TIN2 gene has a gene structure similar to that of B6 (data not shown). A TIN2 gene targeting vector was constructed as shown in Fig. 1A. A neomycin resistance gene was inserted into the first exon of TIN2 upstream of the ATG translation initiation codon. To screen for targeting of the vector to the TIN2 locus in ES cells, ES cell DNA was digested with ApaI and blotted with probe A as shown in Fig. 1B. The 8.5-kb DNA fragment was the TIN2 germ line allele, and the 13-kb fragment represented the targeted allele. Eleven targeted ES cells were identified in 168 clones screened. Two targeted ES clones were used to make chimeric mice. Heterozygous TIN2^{+/-} mice were identified by Southern blot analysis or PCR from the offspring of chimeric mice (Fig. 1B and C). The 220-bp PCR product corresponded to the wild-type allele, and the 270-bp product corresponded to the targeted allele of TIN2 (Fig. 1C).

No homozygous TIN2^{-/-} mice were obtained among 237 offspring of intercrosses of TIN2^{+/-} mice as shown in Table 1. This finding indicated that TIN2 is essential for mouse development and that homozygous inactivation of TIN2 is therefore lethal. The ratio of TIN2^{+/-} to TIN2^{+/+} offspring was approximately 2 to 1, suggesting that there is no haploinsufficiency and decreased viability in TIN2^{+/-} mice.

To determine the time of death during development of TIN2^{-/-} embryos, embryos were isolated from TIN2 heterozygous intercrosses at embryonic day 7.5 and 10.5, and individual embryos were genotyped by PCR or Southern blot analysis. No homozygous TIN2-deficient embryos were found in either day 7.5 litters or day 10.5 litters (Table 2). Thus, TIN2-deficient embryos die before day 7.5 of their embryonic development.

TABLE 2. Day 3.5, but not day 7.5 or 10, TIN2^{-/-} embryos were found from TIN2^{+/-} crosses **AQ: E**

Embryonic day	No. of embryos		
	TIN2 ^{+/+}	TIN2 ^{+/-}	TIN2 ^{-/-}
10.5	7	17	0
7.5	9	15	0
13.5	7	22	9




FIG. 2. (A) PCR genotyping of 3.5-day-old *TIN2* embryos. The 220- and 270-bp PCR products represent the wild-type and targeted alleles, respectively. (B) In vitro-cultured 3.5-day-old embryo in ES cell medium. Pictures in the upper panel represent individual *TIN2*^{+/-} embryos after 5 days of culture. Pictures in the lower panel represent individual *TIN2*^{-/-} embryos after 5 days of culture.

ES cells have a growth and/or survival defect during in vitro culture of *TIN2*-deficient blastocysts. ES cells can be derived from blastocysts, and *TIN2*-deficient ES cell lines could thus be established for further study of *TIN2* function during the earliest stages of embryonic development. Day 3.5 embryos were isolated from *TIN2*^{+/-} intercrosses and cultured in ES cell culture medium. All embryos were of similar appearance and viability at the initiation of culture. Cells were collected and genotyped by PCR after 5 days of culture (Fig. 2A). Approximately one-fourth of ES cell cultures were identified as homozygous *TIN2*^{-/-}, indicating that homozygous knockouts survive to day 3.5 in the expected proportions (Table 2). However, it was striking that *TIN2*^{-/-} cultures were uniformly defective in their survival and differentiation in comparison to wild-type or *TIN2*^{+/-} ES cells. Whereas wild-type ES cell clones grew to form multilayered cell masses, *TIN2*^{-/-} ES cell cultures were flat and cell numbers were much lower than those of wild-type or *TIN2*^{+/-} clones (Fig. 2B). These differences in culture were observed in a masked fashion, since genotype was determined only after completion of culture, and were consistent for 9 *TIN2*^{-/-} cultures and 29 *TIN2*^{+/-} or *TIN2*^{+/+} cultures. A growth and/or survival defect was thus apparent in *TIN2*^{-/-} cells at an early stage of ES cell culture. It should be noted that *TIN2* protein may be essential even in the first days of embryogenesis, where this requirement could be obscured by the persistence of maternal egg-derived mRNA or protein.

Embryonic lethality of *TIN2* deficiency is not telomerase dependent. The known function of *TIN2* involves its interaction with TRF1 to down-regulate the telomerase-dependent elongation of telomeres. It therefore appeared possible that embryonic lethality in *TIN2* knockout mice may be mediated by telomerase-dependent effects. To explore this possibility, *TIN2*^{+/-} mice were bred to mTERT-deficient mice that lack telomerase activity. These telomerase-deficient mice have nor-

TABLE 3. There are no *TIN2*^{-/-} offspring from *TIN2*^{+/-} *TERT*^{-/-} crosses

Growth stage	No. of <i>TIN2</i> ^{+/-} <i>TERT</i> ^{-/-} × <i>TIN2</i> ^{+/-} <i>TERT</i> ^{-/-} offspring		
	<i>TIN2</i> ^{+/+}	<i>TIN2</i> ^{+/-}	<i>TIN2</i> ^{-/-}
G ₁	8	18	0
G ₅	19	31	0

mal embryonic development during their early generations and manifest the effects of telomere dysfunction only after progressive telomere shortening occurs through the breeding of multiple generations with homozygous telomerase deficiency. Intercrosses were carried out between *TIN2*^{+/-} *TERT*^{-/-} mice which were either early (G₁) or late (G₅)-generation mTERT homozygous knockouts. As shown in Table 3, neither *TIN2*^{-/-} *TERT*^{-/-} (G₁) nor *TIN2*^{-/-} *TERT*^{-/-} (G₅) offspring were observed. Thus, the embryonic lethality of *TIN2* deficiency occurs in early and late-generation telomerase-deficient genetic backgrounds, indicating that the requirement for *TIN2* in mouse development is telomerase independent.

DISCUSSION

TIN2 has been identified as a TRF1 binding protein that is essential for the negative regulatory function of TRF1 in modulating telomerase-dependent telomere elongation (12). In the present study, a genetic approach was used to examine the in vivo function of *TIN2*. We found that *TIN2* deficiency in mice resulted in early embryonic lethality in genetic backgrounds with either intact or absent telomerase activity. These findings indicate that *TIN2* is essential in embryonic development and demonstrate a previously undescribed telomerase-independent function of *TIN2*.

TIN2 and TRF1 appear to act as a complex to mediate telomere length regulation. TRF1 interacts with *TIN2* through the TRF1 dimerization domain and the TRF1-binding region of *TIN2* (11). Overexpression of either TRF1 or *TIN2* leads to telomere shortening, whereas expression of either DN TRF1 or DN *TIN2* results in telomerase-dependent telomere elongation (12). It has been proposed that *TIN2* functions in this role by binding to TRF1 dimers and altering the conformation of TRF1 (11). This alteration favors the association of TRF1 with telomeric DNA repeats, which in turn interferes with the accessibility of telomerase to telomere ends. It may therefore be predicted that inactivation of either TRF1 or *TIN2* would result in elongation of telomeres in vivo. To test this prediction, *TIN2* knockout mice were generated in the present study. However, *TIN2* inactivation resulted in early embryonic lethality during mouse development, precluding analysis of the effect of *TIN2* on telomere length regulation. Similarly, the rapid death of *TIN2*^{-/-} ES cells in culture has prevented analysis of the mechanism mediating lethality. Recently, Karlseder et al. (10) reported that inactivation of the mouse TRF1 gene results in embryonic lethality and that TRF1 knockout blastocysts have a cell growth defect and increased apoptosis. The phenotype of *TIN2* knockout mice thus appears to be similar to that of TRF1-deficient mice. To understand the roles of *TIN2* and TRF1 in embryonic development and in adult animals, studies

of inducible TIN2 or TRF1 conditional knockout mice will be informative. Inducible inactivation of TIN2 or TRF1 will allow determination of whether defects in telomere length regulation, apoptosis, or cell cycle progression result from gene inactivation and contribute to embryonic lethality in TIN2-deficient mice. We have, in fact, generated TIN2 conditional knockout constructs by using *cre/loxP* techniques and will use these constructs in studies of inducible and tissue-specific TIN2 inactivation.

Our studies have indicated that telomerase inactivation does not reverse the embryonic lethality of TIN2 deficiency. This demonstrates that TIN2 has a telomerase-independent function that is required during mouse development. These observations imply that, in addition to the telomerase-dependent functions played by TIN2-TRF1 complexes, both TIN2 and TRF1 also function in telomerase-independent roles. Additional telomere-associated proteins may be involved in the telomerase-independent function of TIN2 and TRF1. Among these, constitutive knockouts of some proteins such as ku and ATM are not lethal and are therefore not straightforward candidates for mediation of TIN2 or TRF1 lethality. Others, such as tankyrase 1 and 2 are among the candidates that merit further study.

ACKNOWLEDGMENTS

We are grateful to Nanping Weng and Karen S. Hathcock for critical reading of the manuscript. We thank Eileen Southon and Susan Reid for help in generating mutant mice. We thank Genevieve Sanchez-Howard and the staff at Bioqual for expert animal care and breeding.

REFERENCES

- Baumann, P., and T. R. Cech. 2001. Pot1, the putative telomere end-binding protein in fission yeast and humans. *Science* **292**:1171–1175.
- Bianchi, A., S. Smith, L. Chong, P. Elias, and T. de Lange. 1997. TRF1 is a dimer and bends telomeric DNA. *EMBO J.* **16**:1785–1794.
- Bonin, A., S. W. Reid, and L. Tessarollo. 2001. Gene knockouts: isolation, microinjection and transfer of mouse blastocysts. *Methods Mol. Biol.* **158**:121–134.
- Broccoli, D., L. Chong, S. Oelmann, A. A. Fernald, N. Marziliano, B. van Steensel, D. Kipling, M. M. Le Beau, and T. de Lange. 1997. Comparison of the human and mouse genes encoding the telomeric protein, TRF1: chromosomal localization, expression and conserved protein domains. *Hum. Mol. Genet.* **6**:69–76.
- Broccoli, D., A. Smogorzewska, L. Chong, and T. de Lange. 1997. Human telomeres contain two distinct Myb-related proteins, TRF1 and TRF2. *Nat. Genet.* **17**:231–235.
- Bryan, T. M., and T. R. Cech. 1999. Telomerase and the maintenance of chromosome ends. *Curr. Opin. Cell Biol.* **11**:318–324.
- Espejel, S., S. Franco, A. Sgura, D. Gae, S. M. Bailey, G. E. Taccioli, and M. A. Blasco. 2002. Functional interaction between DNA-PKcs and telomerase in telomere length maintenance. *EMBO J.* **21**:6275–6287.
- Hsu, H. L., D. Gilley, S. A. Galande, M. P. Hande, B. Allen, S. H. Kim, G. C. Li, J. Campisi, T. Kohwi-Shigematsu, and D. J. Chen. 2000. Ku acts in a unique way at the mammalian telomere to prevent end joining. *Genes Dev.* **14**:2807–2812.
- Kaminker, P. G., S. H. Kim, R. D. Taylor, Y. Zebardjian, W. D. Funk, G. B. Morin, P. Yaswen, and J. Campisi. 2001. TANK2, a new TRF1-associated poly(ADP-ribose) polymerase, causes rapid induction of cell death upon overexpression. *J. Biol. Chem.* **276**:35891–35899.
- Karlseder, J., L. Kachatrian, H. Takai, K. Mercer, S. Hingorani, T. Jacks, and T. de Lange. 2003. Targeted deletion reveals an essential function for the telomere length regulator Trf1. *Mol. Cell. Biol.* **23**:6533–6541.
- Kim, S. H., S. Han, Y. H. You, D. J. Chen, and J. Campisi. 2003. The human telomere-associated protein TIN2 stimulates interactions between telomeric DNA tracts in vitro. *EMBO Rep.* **4**:685–691.
- Kim, S. H., P. Kaminker, and J. Campisi. 1999. TIN2, a new regulator of telomere length in human cells. *Nat. Genet.* **23**:405–412.
- Kishi, S., X. Z. Zhou, Y. Ziv, C. Khoo, D. E. Hill, Y. Shiloh, and K. P. Lu. 2001. Telomeric protein Pin2/TRF1 as an important ATM target in response to double strand DNA breaks. *J. Biol. Chem.* **276**:29282–29291.
- Li, B., S. Oestreich, and T. de Lange. 2000. Identification of human Rap1: implications for telomere evolution. *Cell* **101**:471–483.
- Loayza, D., and T. De Lange. 2003. POT1 as a terminal transducer of TRF1 telomere length control. *Nature* **424**:1013–1018.
- Martin, G. R. 1981. Isolation of a pluripotent cell line from early mouse embryos cultured in medium conditioned by teratocarcinoma stem cells. *Proc. Natl. Acad. Sci. USA* **78**:7634–7638.
- Smith, A. G., and M. L. Hooper. 1987. Buffalo rat liver cells produce a diffusible activity which inhibits the differentiation of murine embryonal carcinoma and embryonic stem cells. *Dev. Biol.* **121**:1–9.
- Smith, S., and T. de Lange. 1997. TRF1, a mammalian telomeric protein. *Trends Genet.* **13**:21–26.
- Smith, S., I. Giriat, A. Schmitt, and T. de Lange. 1998. Tankyrase, a poly(ADP-ribose) polymerase at human telomeres. *Science* **282**:1484–1487.
- Tessarollo, L. 2001. Gene knockouts: manipulating mouse embryonic stem cells. *Methods Mol. Biol.* **158**:47–63.
- van Steensel, B., and T. de Lange. 1997. Control of telomere length by the human telomeric protein TRF1. *Nature* **385**:740–743.
- Zhou, X. Z., and K. P. Lu. 2001. The Pin2/TRF1-interacting protein PinX1 is a potent telomerase inhibitor. *Cell* **107**:347–359.

AUTHOR QUERIES

AUTHOR PLEASE ANSWER ALL QUERIES

1

AQA—If FIAU not spelled out as meant, please correct here.

AQB—Please provide initial(s) for Chiang here. Also, please provide initials and surnames for subsequent authors.

AQC—Please give the “x g” value or the brand name of the instrument for centrifugation here.

AQD—Table 1 redrawn. Please check data carefully.

AQE—Table 2 redrawn. Please check data carefully.

AQF—Table 3 redrawn. Please check data carefully.
